



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Feasibility of a multiparametric MRI protocol for imaging biomarkers associated with neoadjuvant radiotherapy for soft tissue sarcoma

Citation for published version:

Kershaw, L, Forker, L, Roberts, D, Sanderson, B, Shenjere, P, Wylie, J, Coyle, C, Kochhar, R, Manoharan, P & Choudhury, A 2021, 'Feasibility of a multiparametric MRI protocol for imaging biomarkers associated with neoadjuvant radiotherapy for soft tissue sarcoma', *BJR|Open*, vol. 3, no. 1, 20200061.
<https://doi.org/10.1259/bjro.20200061>

Digital Object Identifier (DOI):

[10.1259/bjro.20200061](https://doi.org/10.1259/bjro.20200061)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

BJR|Open

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Feasibility of a multiparametric MRI protocol for imaging biomarkers associated with neoadjuvant radiotherapy for soft tissue sarcoma --Manuscript Draft--

Manuscript Number:	BJRO-D-20-00061R1
Full Title:	Feasibility of a multiparametric MRI protocol for imaging biomarkers associated with neoadjuvant radiotherapy for soft tissue sarcoma
Short Title:	Feasibility of mpMRI evaluation of radiotherapy response in STS
Article Type:	Original research
Keywords:	Soft Tissue Sarcoma; Neoadjuvant radiotherapy; MRI; Diffusion weighted imaging; Dynamic contrast-enhanced imaging; Hypoxia
Corresponding Author:	Lucy Kershaw The University of Edinburgh Edinburgh, Edinburgh City UNITED KINGDOM
Corresponding Author Secondary Information:	
Corresponding Author's Institution:	The University of Edinburgh
Corresponding Author's Secondary Institution:	
First Author:	Lucy Kershaw
First Author Secondary Information:	
Order of Authors:	Lucy Kershaw
	Laura Forker
	Darren Roberts
	Benjamin Sanderson
	Patrick Shenjere
	James Wylie
	Catherine Coyle
	Rohit Kochhar
	Ananaya Choudhury
	Prakash Manoharan
Order of Authors Secondary Information:	
Abstract:	<p>Objectives Soft tissue sarcoma (STS) is a rare malignancy with a 5 year overall survival rate of 55%. Neoadjuvant radiotherapy is commonly used in preparation for surgery, but methods to assess early response are lacking despite pathological response at surgery being predictive of overall survival, local recurrence and distant metastasis. Multiparametric MR imaging (mpMRI) is used to assess response in a variety of tumours but lacks a robust, standardised method. The overall aim of this study was to develop a feasible imaging protocol to identify imaging biomarkers for further investigation. Methods Fifteen patients with biopsy-confirmed STS suitable for preoperative radiotherapy and radical surgery were imaged throughout treatment. The mpMRI protocol included anatomical, diffusion weighted and dynamic contrast-enhanced imaging, giving estimates of apparent diffusion coefficient (ADC) and the area under the enhancement curve at 60 s (iAUC 60). Histological analysis of resected tumours included detection of CD31, Ki67, HIF and calculation of a hypoxia score. Results There was a significant reduction in T1 at visit 2 and in ADC at visit 3. Significant associations were found between hypoxia and pre-treatment iAUC 60 , pre-treatment ADC and mid-treatment iAUC 60 . There was also statistically significant</p>

	association between mid-treatment ADC and Ki67. Conclusions This work showed that mpMRI throughout treatment is feasible in patients with STS having neoadjuvant radiotherapy. The relationships between imaging parameters, tissue biomarkers and clinical outcomes warrant further investigation. Advances in Knowledge mpMRI based biomarkers have good correlation with STS tumour biology and are potentially of use for evaluation of radiotherapy response.
--	--

Response to Reviewer 2

1- Missing Data

The expected sample size was 15 patients and the available data was less than that. If the study considered completed, is it safe to use missing data? If the collected data was enough for the authors before reaching 15 patients, do we consider this result as a preliminary/interim result? Is the interim analysis planned? Does that mention in the protocol? If not, the authors should include them as an amendment.

The study design was for a sample size of 15 patients, which was felt to be adequate to assess the feasibility of applying this protocol in this patient group. We are not planning to enrol any additional patients and we consider the study to be complete. Our apologies, we are unsure exactly what the reviewer means by 'is it safe to use missing data?'. We included all the collected data in our analysis and we believe that this is appropriate for detecting groupwise changes. We have added a line in the results section to emphasise this:

Briefly, by the close of the study 15 patients were recruited and twelve scanned (nine had three scans, three had only the first two). All collected data were included in our analysis.

I found reluctance in presentations of results in the discussion part, such as the following statements:

- i). Incomplete datasets were due to incorrect protocols being used, rather than poor patient compliance*
- ii) Postradiotherapy the correlation was not significant, but there were only ten observations at this stage.*

We have reworded the first point as:

- i) The imaging protocol is deliverable as shown by the good patient compliance, and there are some interesting findings

Point 2 was removed.

2- The discussion part was very short.

My suggestion to correlate positive findings (such as iAUC60 and hypoxia score and mid-treatment ADC and Ki67) in this study for specific histopathology, soft tissue sarcoma, with different tumors, in clinical and Preclinical settings of other studies.

Many thanks for this suggestion. We have added the following text and references to the discussion:

AUC inversely correlated with hypoxia score

This is consistent with previous work in a mouse xenograft model, which showed reduced AUC in hypoxic tumour regions defined by pimonidazole staining²⁹

Mid-Tx ADC inversely correlated with Ki67 at resection

The relationship between ADC and Ki67 has been explored previously, and reported in a meta-analysis that confirmed this negative correlation in many tumour types.

No CAIX at surgery have higher ADC mid Tx

In a previous study in melanoma xenografts, ADC was shown to be inversely related to hypoxic fraction determined by pimonidazole staining³¹. The relationship between CAIX and ADC has been explored previously but no relationship was found³², possibly because (unlike pimonidazole staining) CAIX is a downstream marker of hypoxia, influenced by other factors.

Title page

Title: Feasibility of a multiparametric MRI protocol for imaging biomarkers associated with neoadjuvant radiotherapy for soft tissue sarcoma

Short title: Feasibility of mpMRI evaluation of radiotherapy response in STS

Manuscript type: Full paper

Authors:

Lucy E Kershaw PhD, The University of Manchester, Manchester Academic Health Science Centre, The Christie NHSFT, Wilmslow Road, Manchester M20 4BX

Laura Forker MBChB Hons MRes, Translational Radiobiology Group, Division of Cancer Sciences, The University of Manchester, Manchester Academic Health Science Centre, The Christie NHSFT, Wilmslow Road, Manchester M20 4BX

Darren Roberts PhD, Translational Radiobiology Group, Division of Cancer Sciences, The University of Manchester, Manchester Academic Health Science Centre, The Christie NHSFT, Wilmslow Road, Manchester M20 4BX

Benjamin Sanderson MBBS, The University of Manchester, Manchester Academic Health Science Centre, The Christie NHSFT, Wilmslow Road, Manchester M20 4BX

Patrick Shenjere MBChB, MMed(Histopath), FRCPath, FEBP, Dept of Histopathology, The Christie NHSFT, Wilmslow Road, Manchester M20 4BX

James Wylie MBBS, MRCP, FRCR, Dept of Clinical Oncology, The Christie NHSFT, Wilmslow Road, Manchester M20 4BX

Catherine Coyle MB, Bch BaO, MRCP, FRCR, PGCE, Dept of Clinical Oncology, The Christie NHSFT, Wilmslow Road, Manchester M20 4BX

Rohit Kochhar MBBS, MD, DNB, FRCR, Dept of Radiology, The Christie NHSFT, Wilmslow Road, Manchester M20 4BX

Joint last authors:

Ananya Choudhury MA, PhD, MRCP, FRCR, Translational Radiobiology Group, Division of Cancer Sciences, The University of Manchester, Manchester Academic Health Science Centre, The Christie NHSFT, Wilmslow Road, Manchester M20 4BX

Prakash Manoharan MBChB MRCP FRCR, Dept of Radiology, The Christie NHSFT, Wilmslow Road, Manchester M20 4BX

Conflicts of interest and funding statement

The authors have no conflicts of interest to declare. Professor Ananya Choudhury is supported by the NIHR Manchester Biomedical Research Centre. This research was supported by CTRad.

Acknowledgements

We acknowledge the help of Sarah Welby, radiographers at The Christie NHSFT and Prof Catharine West.

Abstract

Objectives

Soft tissue sarcoma (STS) is a rare malignancy with a 5 year overall survival rate of 55%. Neoadjuvant radiotherapy is commonly used in preparation for surgery, but methods to assess early response are lacking despite pathological response at surgery being predictive of overall survival, local recurrence and distant metastasis. Multiparametric MR imaging (mpMRI) is used to assess response in a variety of tumours but lacks a robust, standardised method. The overall aim of this study was to develop a feasible imaging protocol to identify imaging biomarkers for further investigation.

Methods

Fifteen patients with biopsy-confirmed STS suitable for preoperative radiotherapy and radical surgery were imaged throughout treatment. The mpMRI protocol included anatomical, diffusion weighted and dynamic contrast-enhanced imaging, giving estimates of apparent diffusion coefficient (ADC) and the area under the enhancement curve at 60 s (iAUC₆₀). Histological analysis of resected tumours included detection of CD31, Ki67, HIF and calculation of a hypoxia score.

Results

There was a significant reduction in T1 at visit 2 and in ADC at visit 3. Significant associations were found between hypoxia and pre-treatment iAUC₆₀, pre-treatment ADC and mid-treatment iAUC₆₀. There was also statistically significant association between mid-treatment ADC and Ki67.

Conclusions

This work showed that mpMRI throughout treatment is feasible in patients with STS having neoadjuvant radiotherapy. The relationships between imaging parameters, tissue biomarkers and clinical outcomes warrant further investigation.

Advances in Knowledge

mpMRI based biomarkers have good correlation with STS tumour biology and are potentially of use for evaluation of radiotherapy response.

Blind Title page

Title: Feasibility of a multiparametric MRI protocol for imaging biomarkers associated with neoadjuvant radiotherapy for soft tissue sarcoma

Short title: Feasibility of mpMRI evaluation of radiotherapy response in STS

Abstract

Objectives

Soft tissue sarcoma (STS) is a rare malignancy with a 5 year overall survival rate of 55%. Neoadjuvant radiotherapy is commonly used in preparation for surgery, but methods to assess early response are lacking despite pathological response at surgery being predictive of overall survival, local recurrence and distant metastasis. Multiparametric MR imaging (mpMRI) is used to assess response in a variety of tumours but lacks a robust, standardised method. The overall aim of this study was to develop a feasible imaging protocol to identify imaging biomarkers for further investigation.

Methods

Fifteen patients with biopsy-confirmed STS suitable for preoperative radiotherapy and radical surgery were imaged throughout treatment. The mpMRI protocol included anatomical, diffusion weighted and dynamic contrast-enhanced imaging, giving estimates of apparent diffusion coefficient (ADC) and the area under the enhancement curve at 60 s (iAUC₆₀). Histological analysis of resected tumours included detection of CD31, Ki67, HIF and calculation of a hypoxia score.

Results

There was a significant reduction in T1 at visit 2 and in ADC at visit 3. Significant associations were found between hypoxia and pre-treatment iAUC₆₀, pre-treatment ADC and mid-treatment iAUC₆₀. There was also statistically significant association between mid-treatment ADC and Ki67.

Conclusions

This work showed that mpMRI throughout treatment is feasible in patients with STS having neoadjuvant radiotherapy. The relationships between imaging parameters, tissue biomarkers and clinical outcomes warrant further investigation.

Advances in Knowledge

mpMRI based biomarkers have good correlation with STS tumour biology and are potentially of use for evaluation of radiotherapy response.

Introduction

Soft tissue sarcoma (STS) is a rare and heterogeneous malignancy. There were 3298 new diagnoses in the UK in 2010, and the 5-year overall survival rate was 55%¹. Tumour grade is prognostic, with 5-year metastasis-free survival rates of 71% and 44% for intermediate and high grade tumours respectively². Surgery, the main treatment for STS of the limb and trunk, involves excising the lesion along with a tumour-free margin of around 1-2 cm, subject to anatomical constraints. In intermediate/high grade tumours or those which require extensive surgery, this is often not possible without risk to function. In these situations, radiotherapy is employed either pre- or post-operatively. Post-operative radiotherapy requires a higher dose being delivered to a larger volume but can be reserved for situations where an inadequate margin is obtained. Pre-operative radiotherapy may be used to shrink a tumour, for some subtypes³ making a tumour more operable⁴, or to sterilise the margins in preparation for excision⁵. The use of a lower dose and reduction in the irradiated volume for pre-operative compared to post-operative radiotherapy results in similar local control, but with a reduction in late toxicity and better function⁶. There is no consensus on the optimal approach for radiotherapy in STS, with a rationale for both pre and post-surgical radiotherapy.

There is currently no method to assess early response to pre-operative radiotherapy, despite pathological response at surgery being predictive of overall survival, local recurrence⁷ and distant metastasis^{8,9}. Imaging biomarkers are attractive because they are non-invasive, and MR is available in most centres. Volume change in STS during pre-operative radiotherapy is minimal despite marked pathological response³, and tumour size changes (RECIST criteria and/or three-dimensional tumour volumes) are poor predictors of tumour-free surgical margins, local control⁴ and overall survival¹⁰.

Multiparametric MR imaging (mpMRI) has been used to demonstrate radiotherapy response early in treatment, which might allow prompt progression to definitive treatment in poorly-responding tumours^{11–13}. Conversely, some STS such as myxoid liposarcoma may respond adequately with lower doses of radiotherapy reducing late effects without compromising outcome¹⁴. Use of functional measures such as dynamic contrast-enhanced (DCE) MRI (measuring tissue microvasculature), and diffusion-weighted imaging (DWI) (measuring restriction of water molecule diffusion) has been shown to increase the ability of imaging to reflect the amount of tumour necrosis^{15–18}, but protocols varied widely and image analysis often required careful selection of tumour subregions. Delivery of a robust protocol suitable for large multicentre studies is challenging. This study aligns with domain I of the imaging biomarker road map¹⁹, assessing the feasibility of delivering a robust protocol across different anatomical sites, but within a single institute.

The overall aim of this study was to identify imaging parameters suitable for investigation as prognostic factors in a larger study. The specific aims were to (i) develop a well-tolerated imaging protocol allowing for repeat scanning, (ii) identify imaging parameters that change significantly during radiotherapy, and (iii) determine relationships between imaging parameters and histological features in resected tumour tissue. As part of this exploratory assessment, particular attention was focussed on post-surgical hypoxic regions in the resected tumour, a parameter known to affect survival^{20,21}.

Materials and Methods

Patients

In this prospective study, fifteen patients with biopsy-confirmed intermediate or high-grade STS suitable for preoperative radiotherapy (50 Gy in 25 fractions) and radical surgery were recruited. This study had a favourable ethical opinion (13/NW/0500) and all patients gave written informed consent. Patient characteristics are summarised in Table 1.

Patients underwent MRI before radiotherapy (within 4 weeks of the start of treatment), in week 2 or 3 of radiotherapy, and 2-4 weeks after radiotherapy. Within one month of the post-radiotherapy visit, the tumours were surgically resected. At surgery, proximal, distal, medial, lateral, depth and superficial margins were demarcated.

Imaging protocol

Patients were imaged at 1.5 T (Siemens Magnetom Avanto, Siemens Healthineers, Erlangen, Germany) using either the peripheral or body matrix coil with the appropriate elements of the spine array. Imaging began with standard clinical sequences (pre-contrast transverse and coronal T1w and T2w, post-contrast transverse and coronal T1w with fat saturation, all turbo spin-echo) and continued with trial sequences as shown in table 2. During the dynamic sequence, 0.1 ml/kg Gadovist was injected using a power injector at 2 ml/s, followed by 20 ml saline at the same rate.

Image Analysis

ADC maps were calculated at the time of acquisition. Images were analysed using Python (Version 3.6). T1 maps were calculated by fitting the saturation recovery turbo FLASH equation on a pixel-by-pixel basis²². The integrated area under the curve in the first 60 seconds after injection ($iAUC_{60}$) was calculated on a pixel-by-pixel basis using trapezoidal integration, after converting the signal intensity vs time curves to contrast agent concentration vs time curves²³.

Tumours were outlined on T2w images by BS (confirmed by a consultant radiologist with expertise in MRI), and volumes automatically calculated from the known voxel size. These regions of interest (ROIs) were transferred to the dynamic images by nearest-neighbour interpolation and to the ADC maps by referring to anatomical landmarks to ensure that the tumour was correctly outlined even in the presence of distortion. ROIs were eroded in-

plane by one pixel to avoid partial volume effects at the region edges. The median and interquartile range over the whole tumour ROI was calculated for ADC, iAUC₆₀ and T1.

Histological Analysis of Resected Tumours

Immunohistochemistry was performed on 4 µm sections from formalin-fixed paraffin-embedded (FFPE) tumour resection samples to score hypoxia inducible factor – 1 alpha (HIF-1 α), carbonic anhydrase IX (CAIX), antigen KI-67 (Ki67) and CD31. HIF-1 α, CAIX and Ki67 staining was performed using the Bond-Max Automated staining system (Leica Biosystems, Milton Keynes, UK). Slides were de-waxed and rehydrated and antigen retrieval was carried out at pH 9.0 for 40 min at 100° C. The primary antibodies were HIF-1a (BD Biosciences 610959; 1:100 dilution), CAIX (NCL-L-CAIX, Novascstra, Leica Biosystems; 1:100 dilution), Ki67 (clone MIB-1, Dako M7240; 1:100 dilution), and CD31 (M0823 Dako; 1:50 dilution). For HIF-1 α, Ki67 and CD31 the negative control was mouse IgG1 (Dako X0931) and for CAIX was mouse IgG2a (Dako, Ely, UK, X0943). All dilutions were in antibody diluent (Leica AR9352) and negative controls were diluted to the same protein concentration as the primary. Slides were stained using a standard BOND processing protocol (available on request) and Bond Polymer Refine Detection System (Leica DS9800). Colorectal cancer cell line spheroids with a diameter <500 µm were used as a positive control for HIF-1 α and CAIX. A FFPE biopsy of normal human placenta was used as a positive control for CD31.

The percentage of tumour cells per core expressing membranous CAIX was scored by a sarcoma pathologist (PS) at ×8 magnification with negative controls available for comparison. Other markers were scored using automated image analysis (Definiens tissue studio v.4.2; Definiens, Munich, Germany).

The percentage of tumour material was assessed by a sarcoma pathologist (PS) on a separate H&E stained section. RNA from three 10µm sections was extracted using the FFPE

RNA/DNA Purification Plus Kit (Norgen, Thorold, Ontario, Canada) including DNase I treatment. The High-Capacity cDNA Reverse Transcription Kit (Life Technologies, Paisley, UK) was used to reverse transcribe total RNA. cDNA was preamplified using a custom pool of TaqMan assays (Life Technologies) and TaqMan PreAmp Master Mix (Life Technologies).

Expression of a 24-gene hypoxia signature derived for STS²⁴ and two endogenous control genes selected for STS²⁵ was determined using custom 384-well TaqMan array cards (Life Technologies) on a QuantStudio 12K Flex Real-Time PCR System (Life Technologies) using TaqMan Fast Advanced Master Mix (Life Technologies) according to the manufacturer's guidelines. The geometric mean of the endogenous control genes was used for normalisation. Hypoxia scores (HS) were calculated as the normalised median expression of the 24 hypoxia genes (note that the median is used due to the small sample size in this study, rather than the method presented previously²⁴).

Statistical Analysis

In this small feasibility study, changes in median T1, ADC and iAUC₆₀ between the three visits were assessed using a Wilcoxon signed ranks test. Correlations between imaging and histology parameters were assessed by calculating the Pearson correlation coefficient and its associated p-value using the t distribution. Differences in imaging parameters between the tumours with and without CAIX staining was assessed using the Mann-Whitney U test. No correction was made for multiple comparisons.

Results

Patients

Figure 1 shows a flow diagram for enrolment, imaging and analysis. Briefly, by the close of the study 15 patients were recruited and twelve scanned (nine had three scans, three had only the first two). All collected data were included in our analysis. As of January 2019,

Commented [KL1]: Expanded in response to reviewer 2.

one patient had died from acute myeloid leukaemia and two had developed metastatic disease (one patient had a lung metastasis resected and the other developed a solitary spinal metastasis).

Image Analysis

No significant reduction in volume was observed across visits. The median volumes with their interquartile ranges in cm³ were: 29 (22-51) for visit 1, 34 (23-48) for visit 2 and 25 (10-32) for visit 3. In comparison with pre-radiotherapy values, there was a significant reduction in T1 at visit 2 ($p=0.008$) (Figure 2) and in ADC at visit 3 ($p=0.04$), with example ADC maps shown in Fig 3 for two patients. No significant changes in $iAUC_{60}$ were seen over the three visits.

Histological results

Pearson correlation coefficients between imaging and histological parameters are shown with their p-values in Table 3. Significant correlations were found between hypoxia scores and pre-treatment $iAUC_{60}$ ($r = -0.64$, $p = 0.03$), pre-treatment ADC ($r = 0.63$, $p = 0.03$) and mid-treatment $iAUC_{60}$ ($r = -0.63$, $p = 0.03$). There was also a significant correlation between mid-treatment ADC and Ki67 ($r = -0.66$, $p=0.02$). Stratification of patients by CAIX staining demonstrated significant differences in T1 (visit 1 $p = 0.003$, visit 2 $p = 0.01$, visit 3 $p = 0.03$), and $iAUC_{60}$ at visit 1 ($p = 0.03$).

Discussion

In this work we developed a mpMRI protocol for STS, including established functional and structural imaging, that was acceptable for patients and that could be applied several times during radiotherapy. The imaging protocol is deliverable as shown by the good patient compliance, and there are some interesting findings. To our knowledge no other study has

Commented [KL2]: Changed in response to reviewer
2

1
2
3
4
5
6
7 reported the use of mpMRI in STS in neoadjuvant radiotherapy with a time point early in
8
9 treatment..

10
11
12 We applied the techniques of DCE-MRI, DWI and T1 measurement to explore radiotherapy-
13
14 related changes to tumour tissue not reflected by changes in size. As found in several
15
16 previous publications^{3,4,10}, size change varied between tumours and was not related to any
17
18 histological parameters measured at surgery. Since conventional RECIST criteria cannot be
19
20 associated with radiotherapy response, a non-invasive imaging biomarker predictive of
21
22 overall survival, local control⁷ and distant metastasis^{8,9} is desirable and could be used to
23
24 stratify patients for treatment intensification or de-intensification. The tissue T1 decreased
25
26 significantly between baseline and mid-treatment but by the end of radiotherapy the main
27
28 variation in T1 was between patients. T1 changes can reflect a wide range of alterations in
29
30 tissue structure²⁶ resulting in large variations in values between patients, which complicates
31
32 interpretation of tumour revascularisation between baseline and early treatment. Tumours
33
34 with positive staining for CAIX had a significantly shorter T1 at all three visits, consistent
35
36 with the expected T1 shortening effect of deoxyhaemoglobin. ADC increased significantly
37
38 between baseline and post-treatment, as shown in previous work²⁷ and is hypothesised to
39
40 reflect decreased cellularity and increased necrosis²⁸. Pre-treatment ADC values were similar
41
42 to those reported in previous studies^{27,28} though, as noted by other authors, the range of
43
44 baseline values was large.

45
46 There was an inverse correlation between $iAUC_{60}$ and hypoxia score. $iAUC_{60}$ is a semi-
47
48 quantitative parameter with no direct physiological interpretation. A source of the negative
49
50 correlation observed both at baseline and early in treatment could be poor tumour
51
52 perfusion resulting in a lower $iAUC_{60}$ and the post treatment hypoxia observed. This is
53
54 consistent with previous work in a mouse xenograft model, which showed reduced AUC in
55
56 hypoxic tumour regions defined by pimonidazole staining²⁹. At baseline, tumours with

Commented [KL3]: Added in response to reviewer 2.

1
2
3
4
5
6
7 positive staining for post-treatment CAIX had significantly higher $iAUC_{60}$, which is not
8 consistent with the expected relationship between $iAUC_{60}$, perfusion and oxygenation. The
9 inverse correlation between mid-treatment ADC and Ki67 at resection suggests that tumours
10 with a good initial response to radiotherapy (lower cell density, higher ADC) subsequently
11 have less proliferation at resection. The relationship between ADC and Ki67 has been
12 explored previously, and reported in a meta-analysis that confirmed this negative correlation
13 in many tumour types³⁰. Similarly, tumours that show no staining for CAIX at resection
14 (normoxic at surgery) have a significantly higher ADC early in treatment (lower cell density).
15
16 In a previous study in melanoma xenografts, ADC was shown to be inversely related to
17 hypoxic fraction determined by pimonidazole staining³¹. The relationship between CAIX
18 and ADC has been explored previously but no relationship was found³², possibly because
19 CAIX is a downstream marker of hypoxia which can be regulated by other factors whereas
20 pimonidazole represents a more direct measure. Correlation between pre-treatment ADC
21 and hypoxia score is more difficult to interpret as ADC is measured long before the resection
22 of the tumour.
23
24
25
26
27
28
29
30
31
32

Commented [KL4]: Added in response to reviewer 2

Commented [KL5]: Added in response to reviewer 2

33
34
35 This study has several limitations. The number of patients was small, prohibiting
36 examination of differences between responders and non-responders. The results should be
37 interpreted with caution due to sample size and differences in measurement timepoint, but
38 the main aim of the study was to develop a suitable imaging protocol for which this small
39 number is likely to be sufficient. The DCE-MRI data were acquired with sufficient temporal
40 resolution to support tracer kinetics modelling, but the varying tumour locations made
41 measurement of an arterial input function extremely challenging. We therefore opted to
42 use a semi-quantitative parameter instead, but modelling could perhaps have given further
43 insight¹⁷. Future work could include modelling, if suitable spatial resolution can be obtained,
44 and an MR-linac could allow more detailed monitoring during treatment³³.
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Overall, this work has resulted in a feasible imaging protocol aligning with domain I of the imaging biomarker road map. We identified significant changes in T1 and ADC during treatment. As iAUC relates to hypoxia, an established adverse prognostic factor in STS³⁴, it may be suitable as a non-invasive biomarker of tumour microenvironment and should be explored in a larger study.

References

1. National Cancer Intelligence Network. Bone and Soft Tissue Sarcomas UK Incidence and Survival: 1996 to 2010. *Natl Cancer Intelligence Netw.* 2013;(November 2013):17. <http://ncin.org.uk/view?rid=2353>.
2. Coindre JM, Terrier P, Guillou L, et al. Predictive value of grade for metastasis development in the main histologic types of adult soft tissue sarcomas: A study of 1240 patients from the French Federation of Cancer Centers sarcoma group. *Cancer.* 2001;91(10):1914-1926. doi:10.1002/1097-0142(20010515)91:10<1914::AID-CNCR1214>3.0.CO;2-3
3. Roberge D, Skamene T, Nahal A, Turcotte RE, Powell T, Freeman C. Radiological and pathological response following pre-operative radiotherapy for soft-tissue sarcoma. *Radiother Oncol.* 2010;97(3):404-407. doi:10.1016/j.radonc.2010.10.007
4. Le Grange F, Cassoni AM, Seddon BM. Tumour volume changes following pre-operative radiotherapy in borderline resectable limb and trunk soft tissue sarcoma. *Eur J Surg Oncol.* 2014;40(4):394-401. doi:10.1016/j.ejso.2014.01.011
5. Dangoor A, Seddon B, Gerrand C, Grimer R, Whelan J, Judson I. UK guidelines for the management of soft tissue sarcomas. *Clin Sarcoma Res.* 2016;6(1):20. doi:10.1186/s13569-016-0060-4
6. DAVIS A, OSULLIVAN B, TURCOTTE R, et al. Late radiation morbidity following randomization to preoperative versus postoperative radiotherapy in extremity soft tissue sarcoma. *Radiother Oncol.* 2005;75(1):48-53. doi:10.1016/j.radonc.2004.12.020

7. Eilber FC, Rosen G, Eckardt J, et al. Treatment-Induced Pathologic Necrosis: A Predictor of Local Recurrence and Survival in Patients Receiving Neoadjuvant Therapy for High-Grade Extremity Soft Tissue Sarcomas. Bönig H, ed. *J Clin Oncol*. 2001;19(13):3203-3209. doi:10.1200/JCO.2001.19.13.3203
8. Shah D, Borys D, Martinez SR, et al. Complete pathologic response to neoadjuvant radiotherapy is predictive of oncological outcome in patients with soft tissue sarcoma. *Anticancer Res*. 2012;32(9):3911-3915. doi:10.1038/mp.2011.182.doi
9. MacDermid DM, Miller LL, Peabody TD, et al. Primary Tumor Necrosis Predicts Distant Control in Locally Advanced Soft-Tissue Sarcomas After Preoperative Concurrent Chemoradiotherapy. Bönig H, ed. *Int J Radiat Oncol*. 2010;76(4):1147-1153. doi:10.1016/j.ijrobp.2009.03.015
10. Miki Y, Ngan S, Clark JCMCM, Akiyama T, Choong PFMFM. The significance of size change of soft tissue sarcoma during preoperative radiotherapy. *Eur J Surg Oncol*. 2010;36(7):678-683. doi:10.1016/j.ejso.2010.05.021
11. Pasquier D, Hadj Henni A, Escande A, et al. Diffusion weighted MRI as an early predictor of tumor response to hypofractionated stereotactic boost for prostate cancer. *Sci Rep*. 2018;8(1):10407. doi:10.1038/s41598-018-28817-9
12. Pham TT, Liney GP, Wong K, Barton MB. Functional MRI for quantitative treatment response prediction in locally advanced rectal cancer. *Br J Radiol*. 2017;90(1072):20151078. doi:10.1259/bjr.20151078
13. Suh CH, Kim HS, Jung SC, Choi CG, Kim SJ. Multiparametric MRI as a potential surrogate endpoint for decision-making in early treatment response following concurrent chemoradiotherapy in patients with newly diagnosed glioblastoma: a systematic review and meta-analysis. *Eur Radiol*. 2018;28(6):2628-2638. doi:10.1007/s00330-017-5262-5
14. Chung PWM, Dehesi BM, Ferguson PC, et al. Radiosensitivity translates into excellent local control in extremity myxoid liposarcoma: A comparison with other

- soft tissue sarcomas. *Cancer*. 2009;115(14):3254-3261. doi:10.1002/cncr.24375
15. Soldatos T, Ahlawat S, Montgomery E, Chalian M, Jacobs MA, Fayad LM. Multiparametric Assessment of Treatment Response in High-Grade Soft-Tissue Sarcomas with Anatomic and Functional MR Imaging Sequences. *Radiology*. 2016;278(3):831-840. doi:10.1148/radiol.2015142463
16. Meyer JM, Perlewitz KS, Hayden JB, et al. Phase I Trial of Preoperative Chemoradiation plus Sorafenib for High-Risk Extremity Soft Tissue Sarcomas with Dynamic Contrast-Enhanced MRI Correlates. *Clin Cancer Res*. 2013;19(24):6902-6911. doi:10.1158/1078-0432.CCR-13-1594
17. Xia W, Yan Z, Gao X. Volume fractions of DCE-MRI parameter as early predictor of histologic response in soft tissue sarcoma: A feasibility study. *Eur J Radiol*. 2017;95(July):228-235. doi:10.1016/j.ejrad.2017.08.021
18. Schnapauff D, Zeile M, Niederhagen M Ben, et al. Diffusion-weighted echo-planar magnetic resonance imaging for the assessment of tumor cellularity in patients with soft-tissue sarcomas. *J Magn Reson Imaging*. 2009;29(6):1355-1359. doi:10.1002/jmri.21755
19. O'Connor JPB, Aboagye EO, Adams JE, et al. Imaging biomarker roadmap for cancer studies. *Nat Rev Clin Oncol*. 2017;14(3):169-186. doi:10.1038/nrclinonc.2016.162
20. Maeda-Otsuka S, Kajihara I, Tasaki Y, et al. Hypoxia accelerates the progression of angiosarcoma through the regulation of angiosarcoma cells and tumor microenvironment. *J Dermatol Sci*. 2019. doi:10.1016/j.jdermsci.2019.01.005
21. Forker L, Gaunt P, Sioletic S, et al. The hypoxia marker CAIX is prognostic in the UK phase III Vortex-Biobank cohort: An important resource for translational research in soft tissue sarcoma. *Br J Cancer*. 2018. doi:10.1038/bjc.2017.430
22. Larsson HB, Rosenbaum S, Fritz-Hansen T. Quantification of the effect of water exchange in dynamic contrast MRI perfusion measurements in the brain and heart. *Magn Reson Med*. 2001;46(2):272-281.

- <http://www.ncbi.nlm.nih.gov/pubmed/11477630>.
23. Brookes JA, Redpath TW, Gilbert FJ, Murray AD, Staff RT. Accuracy of T1 measurement in dynamic contrast-enhanced breast MRI using two- and three-dimensional variable flip angle fast low-angle shot. *J Magn Reson Imaging*. 1999;9(2):163-171. <http://www.ncbi.nlm.nih.gov/pubmed/10077009>.
 24. Yang L, Forker L, Irlam JJ, Pillay N, Choudhury A, West CML. Validation of a hypoxia related gene signature in multiple soft tissue sarcoma cohorts. *Oncotarget*. 2018;9(3):3946-3955. doi:10.18632/oncotarget.23280
 25. Betts GNJ, Eustace A, Patiar S, et al. Prospective technical validation and assessment of intra-tumour heterogeneity of a low density array hypoxia gene profile in head and neck squamous cell carcinoma. *Eur J Cancer*. 2013;49(1):156-165. doi:10.1016/j.ejca.2012.07.028
 26. McRobbie DW, Moore EA, Graves MJ, Prince MR. *MRI from Picture to Proton*. Cambridge: CUP; 2003.
 27. Einarsson H, Karlsson M, Wejde J, Bauer HCF. Diffusion-weighted MRI of soft tissue tumours. *Eur Radiol*. 2004;14(6):959-963. doi:10.1007/s00330-004-2237-0
 28. Dudeck O, Zeile M, Pink D, et al. Diffusion-weighted magnetic resonance imaging allows monitoring of anticancer treatment effects in patients with soft-tissue sarcomas. *J Magn Reson Imaging*. 2008;27(5):1109-1113. doi:10.1002/jmri.21358
 29. Jardim-Perassi B V., Huang S, Dominguez-Viqueira W, et al. Multiparametric MRI and coregistered histology identify tumor habitats in breast cancer mouse models. *Cancer Res*. 2019;79(15):3952-3964. doi:10.1158/0008-5472.CAN-19-0213
 30. Surov A, Meyer HJ, Wienke A. Associations between apparent diffusion coefficient (ADC) and KI 67 in different tumors: A meta-analysis. part 2: ADCmin. *Oncotarget*. 2018;9(9):8675-8680. doi:10.18632/oncotarget.24006
 31. Hompland T, Ellingsen C, Galappathi K, Rofstad EK. DW-MRI in assessment of the hypoxic fraction, interstitial fluid pressure, and metastatic propensity of melanoma

1
2
3
4
5
6
7 xenografts. *BMC Cancer*. 2014;14(1):1-9. doi:10.1186/1471-2407-14-92

- 8
9 32. Rasmussen JH, Olin A, Lelkaitis G, et al. Does multiparametric imaging with 18F-FDG-
10 PET/MRI capture spatial variation in immunohistochemical cancer biomarkers in head
11 and neck squamous cell carcinoma? *Br J Cancer*. 2020;123(1):46-53.
12 doi:10.1038/s41416-020-0876-9
13
14 33. Kerkmeijer LGW, Fuller CD, Verkooijen HM, et al. The MRI-Linear Accelerator
15 Consortium: Evidence-Based Clinical Introduction of an Innovation in Radiation
16 Oncology Connecting Researchers, Methodology, Data Collection, Quality Assurance,
17 and Technical Development. *Front Oncol*. 2016;6(October):1-6.
18 doi:10.3389/fonc.2016.00215
19
20 34. Nordsmark M, Alsner J, Keller J, et al. Hypoxia in human soft tissue sarcomas: Adverse
21 impact on survival and no association with p53 mutations. *Br J Cancer*.
22 2001;84(8):1070-1075. doi:10.1054/bjoc.2001.1728
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Tables

Table 1

Patient characteristics

Patient	Age at first scan	M/F	Tumour location	Tumour type, grade, stage	Status as of Jan 2019
1	73	M	Upper arm	Undifferentiated spindle cell G2, T2bN0M0	No disease
2	56	M	Trunk	Myxoid liposarcoma G3, T2bN0M0	No disease
3	79	M	Upper arm	Myxofibrosarcoma G2, T1bN0M0	No disease
4	27	M	Knee	Myxoid liposarcoma G3, T2bN0M0	No disease
5	29	M	Lower leg	Myxoid chondrosarcoma G3, T2bN0M0	No disease
6	69	F	Trunk	Undifferentiated pleomorphic sarcoma G3, T1aN0M0	No disease
7	41	M	Lower leg	Myxofibrosarcoma G3, T2bN0M0	No disease but chronic inflammation post-surgery
8	62	M	Forearm	Myxofibrosarcoma G3, T1bN0M0	No disease
9	24	M	Knee	Synovial sarcoma G3, T2bN0M0	Lung metastasis resected June 2018, now no disease
10	67	M	Knee	Undifferentiated pleomorphic sarcoma G3, T2bN0M0	Died (Acute Myeloid Leukaemia) Feb 2018
11	33	M	Thigh	Myxoid liposarcoma G3, T2bN0M0	Single metastasis in spine August 2018
12	74	M	Trunk	Undifferentiated spindle cell sarcoma G3, T2bN0M0	No disease

Table 2

Trial Imaging protocol

Sequence	Purpose	Flip angle, TE, TR / ms	Parallel imaging factor	Other	Matrix ^a	FOV / cm
TSE 2D Turbo spin- echo	High resolution T2w for tumour outlining	150° 96, 3890	None	ETL 13	256 x 256 x 20	Limb: 25 x 25 x 10 Trunk: 38.6 x 38.6 x 10
EPI 2D Echoplanar imaging	Diffusion- weighted imaging	- 103, 12100	2 AP	EPI factor 128, B = 0, 50, 100, 150, 200, 500, 1000 s/mm ²	128 x 128 x 20	
SRTFE 3D Saturation- recovery turbo field echo	T1 measurement	12° 1.52 64, 142, 292, 1050, 2550, 3950	2 AP	TI = 37, 100, 250, 1000, 2500, 3900 ms		
VIBE 3D Volume- interpolated breath hold imaging	Dynamic contrast- enhanced imaging	16° 0.81, 2.63	2 AP	Temporal resolution 1.75 s, 150 dynamics		

^a In one case, 26 slices were needed to cover the tumour, leading to a dynamic temporal resolution of 3.2 s, and TR values for the SRTFE of 73, 145, 306, 1060, 2560, and 3960 ms.

TE – echo time, TR – repetition time, TI – inversion time, FOV – field of view

Table 3

Pearson correlation coefficients for correlations between imaging and histological parameters., with p-values calculated from a t-distribution shown in brackets.

	T1	iAUC	ADC	Volume
	Pre-treatment			
CD31	-0.07 (0.83)	-0.28 (0.37)	-0.16 (0.63)	-0.37 (0.23)
Ki67	-0.50 (0.10)	0.11 (0.74)	-0.55 (0.06)	-0.29 (0.36)
HIF	0.08 (0.80)	-0.40 (0.19)	0.04 (0.91)	-0.10 (0.76)
Hypoxia score	0.47 (0.12)	-0.64 (0.03)*	0.63 (0.03)*	-0.16 (0.63)
	Mid-treatment			
CD31	-0.09 (0.78)	-0.43 (0.17)	-0.31 (0.32)	-0.32 (0.31)
Ki67	-0.43 (0.16)	-0.05 (0.88)	-0.66 (0.02)*	-0.31 (0.33)
HIF	0.03 (0.94)	-0.44 (0.16)	-0.10 (0.75)	-0.13 (0.68)
Hypoxia score	0.24 (0.45)	-0.63 (0.03)*	0.58 (0.05)	-0.14 (0.66)
	Post-treatment			
CD31	-0.15 (0.63)	-0.26 (0.42)	-0.29 (0.36)	-0.13 (0.69)
Ki67	-0.57 (0.05)	-0.09 (0.78)	-0.44 (0.16)	-0.04 (0.66)
HIF	-0.27 (0.40)	-0.20 (0.53)	-0.36 (0.25)	-0.29 (0.37)
Hypoxia score	0.23 (0.43)	-0.27 (0.40)	0.34 (0.27)	-0.18 (0.57)

Blind Title page

Title: Feasibility of a multiparametric MRI protocol for imaging biomarkers associated with neoadjuvant radiotherapy for soft tissue sarcoma

Short title: Feasibility of mpMRI evaluation of radiotherapy response in STS

Abstract

Objectives

Soft tissue sarcoma (STS) is a rare malignancy with a 5 year overall survival rate of 55%. Neoadjuvant radiotherapy is commonly used in preparation for surgery, but methods to assess early response are lacking despite pathological response at surgery being predictive of overall survival, local recurrence and distant metastasis. Multiparametric MR imaging (mpMRI) is used to assess response in a variety of tumours but lacks a robust, standardised method. The overall aim of this study was to develop a feasible imaging protocol to identify imaging biomarkers for further investigation.

Methods

Fifteen patients with biopsy-confirmed STS suitable for preoperative radiotherapy and radical surgery were imaged throughout treatment. The mpMRI protocol included anatomical, diffusion weighted and dynamic contrast-enhanced imaging, giving estimates of apparent diffusion coefficient (ADC) and the area under the enhancement curve at 60 s (iAUC₆₀). Histological analysis of resected tumours included detection of CD31, Ki67, HIF and calculation of a hypoxia score.

Results

There was a significant reduction in T1 at visit 2 and in ADC at visit 3. Significant associations were found between hypoxia and pre-treatment iAUC₆₀, pre-treatment ADC and mid-treatment iAUC₆₀. There was also statistically significant association between mid-treatment ADC and Ki67.

Conclusions

This work showed that mpMRI throughout treatment is feasible in patients with STS having neoadjuvant radiotherapy. The relationships between imaging parameters, tissue biomarkers and clinical outcomes warrant further investigation.

Advances in Knowledge

mpMRI based biomarkers have good correlation with STS tumour biology and are potentially of use for evaluation of radiotherapy response.

Introduction

Soft tissue sarcoma (STS) is a rare and heterogeneous malignancy. There were 3298 new diagnoses in the UK in 2010, and the 5-year overall survival rate was 55%¹. Tumour grade is prognostic, with 5-year metastasis-free survival rates of 71% and 44% for intermediate and high grade tumours respectively². Surgery, the main treatment for STS of the limb and trunk, involves excising the lesion along with a tumour-free margin of around 1-2 cm, subject to anatomical constraints. In intermediate/high grade tumours or those which require extensive surgery, this is often not possible without risk to function. In these situations, radiotherapy is employed either pre- or post-operatively. Post-operative radiotherapy requires a higher dose being delivered to a larger volume but can be reserved for situations where an inadequate margin is obtained. Pre-operative radiotherapy may be used to shrink a tumour, for some subtypes³ making a tumour more operable⁴, or to sterilise the margins in preparation for excision⁵. The use of a lower dose and reduction in the irradiated volume for pre-operative compared to post-operative radiotherapy results in similar local control, but with a reduction in late toxicity and better function⁶. There is no consensus on the optimal approach for radiotherapy in STS, with a rationale for both pre and post-surgical radiotherapy.

There is currently no method to assess early response to pre-operative radiotherapy, despite pathological response at surgery being predictive of overall survival, local recurrence⁷ and distant metastasis^{8,9}. Imaging biomarkers are attractive because they are non-invasive, and MR is available in most centres. Volume change in STS during pre-operative radiotherapy is minimal despite marked pathological response³, and tumour size changes (RECIST criteria and/or three-dimensional tumour volumes) are poor predictors of tumour-free surgical margins, local control⁴ and overall survival¹⁰.

1 Multiparametric MR imaging (mpMRI) has been used to demonstrate radiotherapy response
2 early in treatment, which might allow prompt progression to definitive treatment in poorly-
3 responding tumours^{11–13}. Conversely, some STS such as myxoid liposarcoma may respond
4 adequately with lower doses of radiotherapy reducing late effects without compromising
5 outcome¹⁴. Use of functional measures such as dynamic contrast-enhanced (DCE) MRI
6 (measuring tissue microvasculature), and diffusion-weighted imaging (DWI) (measuring
7 restriction of water molecule diffusion) has been shown to increase the ability of imaging
8 to reflect the amount of tumour necrosis^{15–18}, but protocols varied widely and image analysis
9 often required careful selection of tumour subregions. Delivery of a robust protocol suitable
10 for large multicentre studies is challenging. This study aligns with domain I of the imaging
11 biomarker road map¹⁹, assessing the feasibility of delivering a robust protocol across
12 different anatomical sites, but within a single institute.
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29

30 The overall aim of this study was to identify imaging parameters suitable for investigation
31 as prognostic factors in a larger study. The specific aims were to (i) develop a well-tolerated
32 imaging protocol allowing for repeat scanning, (ii) identify imaging parameters that change
33 significantly during radiotherapy, and (iii) determine relationships between imaging
34 parameters and histological features in resected tumour tissue. As part of this exploratory
35 assessment, particular attention was focussed on post-surgical hypoxic regions in the
36 resected tumour, a parameter known to affect survival^{20,21}.
37
38
39
40
41
42
43
44
45
46
47

48 **Materials and Methods**

49 **Patients**

50
51 In this prospective study, fifteen patients with biopsy-confirmed intermediate or high-grade
52 STS suitable for preoperative radiotherapy (50 Gy in 25 fractions) and radical surgery were
53 recruited. This study had a favourable ethical opinion (13/NW/0500) and all patients gave
54 written informed consent. Patient characteristics are summarised in Table 1.
55
56
57
58
59
60
61
62
63
64
65

Patients underwent MRI before radiotherapy (within 4 weeks of the start of treatment), in week 2 or 3 of radiotherapy, and 2-4 weeks after radiotherapy. Within one month of the post-radiotherapy visit, the tumours were surgically resected. At surgery, proximal, distal, medial, lateral, depth and superficial margins were demarcated.

Imaging protocol

Patients were imaged at 1.5 T (Siemens Magnetom Avanto, Siemens Healthineers, Erlangen, Germany) using either the peripheral or body matrix coil with the appropriate elements of the spine array. Imaging began with standard clinical sequences (pre-contrast transverse and coronal T1w and T2w, post-contrast transverse and coronal T1w with fat saturation, all turbo spin-echo) and continued with trial sequences as shown in table 2. During the dynamic sequence, 0.1 ml/kg Gadovist was injected using a power injector at 2 ml/s, followed by 20 ml saline at the same rate.

Image Analysis

ADC maps were calculated at the time of acquisition. Images were analysed using Python (Version 3.6). T1 maps were calculated by fitting the saturation recovery turbo FLASH equation on a pixel-by-pixel basis²². The integrated area under the curve in the first 60 seconds after injection (iAUC₆₀) was calculated on a pixel-by-pixel basis using trapezoidal integration, after converting the signal intensity vs time curves to contrast agent concentration vs time curves²³.

Tumours were outlined on T2w images by BS (confirmed by a consultant radiologist with expertise in MRI), and volumes automatically calculated from the known voxel size. These regions of interest (ROIs) were transferred to the dynamic images by nearest-neighbour interpolation and to the ADC maps by referring to anatomical landmarks to ensure that the tumour was correctly outlined even in the presence of distortion. ROIs were eroded in-

plane by one pixel to avoid partial volume effects at the region edges. The median and interquartile range over the whole tumour ROI was calculated for ADC, iAUC₆₀ and T1.

Histological Analysis of Resected Tumours

Immunohistochemistry was performed on 4 µm sections from formalin-fixed paraffin-embedded (FFPE) tumour resection samples to score hypoxia inducible factor – 1 alpha (HIF-1 α), carbonic anhydrase IX (CAIX), antigen KI-67 (Ki67) and CD31. HIF-1 α , CAIX and Ki67 staining was performed using the Bond-Max Automated staining system (Leica Biosystems, Milton Keynes, UK). Slides were de-waxed and rehydrated and antigen retrieval was carried out at pH 9.0 for 40 min at 100° C. The primary antibodies were HIF-1a (BD Biosciences 610959; 1:100 dilution), CAIX (NCL-L-CAIX, Novascastra, Leica Biosystems; 1:100 dilution), Ki67 (clone MIB-1, Dako M7240; 1:100 dilution), and CD31 (M0823 Dako; 1:50 dilution). For HIF-1 α , Ki67 and CD31 the negative control was mouse IgG1 (Dako X0931) and for CAIX was mouse IgG2a (Dako, Ely, UK, X0943). All dilutions were in antibody diluent (Leica AR9352) and negative controls were diluted to the same protein concentration as the primary. Slides were stained using a standard BOND processing protocol (available on request) and Bond Polymer Refine Detection System (Leica DS9800). Colorectal cancer cell line spheroids with a diameter <500 µm were used as a positive control for HIF-1 α and CAIX. A FFPE biopsy of normal human placenta was used as a positive control for CD31.

The percentage of tumour cells per core expressing membranous CAIX was scored by a sarcoma pathologist (PS) at ×8 magnification with negative controls available for comparison. Other markers were scored using automated image analysis (Definiens tissue studio v.4.2; Definiens, Munich, Germany).

The percentage of tumour material was assessed by a sarcoma pathologist (PS) on a separate H&E stained section. RNA from three 10µm sections was extracted using the FFPE

RNA/DNA Purification Plus Kit (Norgen, Thorold, Ontario, Canada) including DNase I treatment. The High-Capacity cDNA Reverse Transcription Kit (Life Technologies, Paisley, UK) was used to reverse transcribe total RNA. cDNA was preamplified using a custom pool of TaqMan assays (Life Technologies) and TaqMan PreAmp Master Mix (Life Technologies).

Expression of a 24-gene hypoxia signature derived for STS²⁴ and two endogenous control genes selected for STS²⁵ was determined using custom 384-well TaqMan array cards (Life Technologies) on a QuantStudio 12K Flex Real-Time PCR System (Life Technologies) using TaqMan Fast Advanced Master Mix (Life Technologies) according to the manufacturer's guidelines. The geometric mean of the endogenous control genes was used for normalisation. Hypoxia scores (HS) were calculated as the normalised median expression of the 24 hypoxia genes (note that the median is used due to the small sample size in this study, rather than the method presented previously²⁴).

Statistical Analysis

In this small feasibility study, changes in median T1, ADC and iAUC₆₀ between the three visits were assessed using a Wilcoxon signed ranks test. Correlations between imaging and histology parameters were assessed by calculating the Pearson correlation coefficient and its associated p-value using the t distribution. Differences in imaging parameters between the tumours with and without CAIX staining was assessed using the Mann-Whitney U test. No correction was made for multiple comparisons.

Results

Patients

Figure 1 shows a flow diagram for enrolment, imaging and analysis. Briefly, by the close of the study 15 patients were recruited and twelve scanned (nine had three scans, three had only the first two). All collected data were included in our analysis. As of January 2019,

one patient had died from acute myeloid leukaemia and two had developed metastatic disease (one patient had a lung metastasis resected and the other developed a solitary spinal metastasis).

Image Analysis

No significant reduction in volume was observed across visits. The median volumes with their interquartile ranges in cm³ were: 29 (22-51) for visit 1, 34 (23-48) for visit 2 and 25 (10-32) for visit 3. In comparison with pre-radiotherapy values, there was a significant reduction in T1 at visit 2 ($p=0.008$) (Figure 2) and in ADC at visit 3 ($p=0.04$), with example ADC maps shown in Fig 3 for two patients. No significant changes in iAUC₆₀ were seen over the three visits.

Histological results

Pearson correlation coefficients between imaging and histological parameters are shown with their p-values in Table 3. Significant correlations were found between hypoxia scores and pre-treatment iAUC₆₀ ($r = -0.64$, $p = 0.03$), pre-treatment ADC ($r = 0.63$, $p = 0.03$) and mid-treatment iAUC₆₀ ($r = -0.63$, $p = 0.03$). There was also a significant correlation between mid-treatment ADC and Ki67 ($r = -0.66$, $p=0.02$). Stratification of patients by CAIX staining demonstrated significant differences in T1 (visit 1 $p = 0.003$, visit 2 $p = 0.01$, visit 3 $p = 0.03$), and iAUC₆₀ at visit 1 ($p = 0.03$).

Discussion

In this work we developed a mpMRI protocol for STS, including established functional and structural imaging, that was acceptable for patients and that could be applied several times during radiotherapy. The imaging protocol is deliverable as shown by the good patient compliance, and there are some interesting findings. To our knowledge no other study has

1 reported the use of mpMRI in STS in neoadjuvant radiotherapy with a time point early in
2 treatment..
3

4
5
6
7 We applied the techniques of DCE-MRI, DWI and T1 measurement to explore radiotherapy-
8 related changes to tumour tissue not reflected by changes in size. As found in several
9 previous publications^{3,4,10}, size change varied between tumours and was not related to any
10 histological parameters measured at surgery. Since conventional RECIST criteria cannot be
11 associated with radiotherapy response, a non-invasive imaging biomarker predictive of
12 overall survival, local control⁷ and distant metastasis^{8,9} is desirable and could be used to
13 stratify patients for treatment intensification or de-intensification. The tissue T1 decreased
14 significantly between baseline and mid-treatment but by the end of radiotherapy the main
15 variation in T1 was between patients. T1 changes can reflect a wide range of alterations in
16 tissue structure²⁶ resulting in large variations in values between patients, which complicates
17 interpretation of tumour revascularisation between baseline and early treatment. Tumours
18 with positive staining for CAIX had a significantly shorter T1 at all three visits, consistent
19 with the expected T1 shortening effect of deoxyhaemoglobin. ADC increased significantly
20 between baseline and post-treatment, as shown in previous work²⁷ and is hypothesised to
21 reflect decreased cellularity and increased necrosis²⁸. Pre-treatment ADC values were similar
22 to those reported in previous studies^{27,28} though, as noted by other authors, the range of
23 baseline values was large.
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

48 There was an inverse correlation between iAUC₆₀ and hypoxia score. iAUC₆₀ is a semi-
49 quantitative parameter with no direct physiological interpretation. A source of the negative
50 correlation observed both at baseline and early in treatment could be poor tumour
51 perfusion resulting in a lower iAUC₆₀ and the post treatment hypoxia observed. This is
52 consistent with previous work in a mouse xenograft model, which showed reduced AUC in
53 hypoxic tumour regions defined by pimonidazole staining²⁹. At baseline, tumours with
54
55
56
57
58
59
60
61
62
63
64
65

1 positive staining for post-treatment CAIX had significantly higher $iAUC_{60}$, which is not
2 consistent with the expected relationship between $iAUC_{60}$, perfusion and oxygenation. The
3
4 inverse correlation between mid-treatment ADC and Ki67 at resection suggests that tumours
5
6 with a good initial response to radiotherapy (lower cell density, higher ADC) subsequently
7
8 have less proliferation at resection. The relationship between ADC and Ki67 has been
9
10 explored previously, and reported in a meta-analysis that confirmed this negative correlation
11
12 in many tumour types³⁰. Similarly, tumours that show no staining for CAIX at resection
13
14 (normoxic at surgery) have a significantly higher ADC early in treatment (lower cell density).
15
16 In a previous study in melanoma xenografts, ADC was shown to be inversely related to
17
18 hypoxic fraction determined by pimonidazole staining³¹. The relationship between CAIX
19
20 and ADC has been explored previously but no relationship was found³², possibly because
21
22 CAIX is a downstream marker of hypoxia which can be regulated by other factors whereas
23
24 pimonidazole represents a more direct measure. Correlation between pre-treatment ADC
25
26 and hypoxia score is more difficult to interpret as ADC is measured long before the resection
27
28 of the tumour.
29
30
31
32
33
34
35

36 This study has several limitations. The number of patients was small, prohibiting
37
38 examination of differences between responders and non-responders. The results should be
39
40 interpreted with caution due to sample size and differences in measurement timepoint, but
41
42 the main aim of the study was to develop a suitable imaging protocol for which this small
43
44 number is likely to be sufficient. The DCE-MRI data were acquired with sufficient temporal
45
46 resolution to support tracer kinetics modelling, but the varying tumour locations made
47
48 measurement of an arterial input function extremely challenging. We therefore opted to
49
50 use a semi-quantitative parameter instead, but modelling could perhaps have given further
51
52 insight¹⁷. Future work could include modelling, if suitable spatial resolution can be obtained,
53
54 and an MR-linac could allow more detailed monitoring during treatment³³.
55
56
57
58
59
60
61
62
63
64
65

Overall, this work has resulted in a feasible imaging protocol aligning with domain I of the imaging biomarker road map. We identified significant changes in T1 and ADC during treatment. As iAUC relates to hypoxia, an established adverse prognostic factor in STS³⁴, it may be suitable as a non-invasive biomarker of tumour microenvironment and should be explored in a larger study.

References

1. National Cancer Intelligence Network. Bone and Soft Tissue Sarcomas UK Incidence and Survival: 1996 to 2010. *Natl Cancer Intelligence Netw.* 2013;(November 2013):17. <http://ncin.org.uk/view?rid=2353>.
2. Coindre JM, Terrier P, Guillou L, et al. Predictive value of grade for metastasis development in the main histologic types of adult soft tissue sarcomas: A study of 1240 patients from the French Federation of Cancer Centers sarcoma group. *Cancer.* 2001;91(10):1914-1926. doi:10.1002/1097-0142(20010515)91:10<1914::AID-CNCR1214>3.0.CO;2-3
3. Roberge D, Skamene T, Nahal A, Turcotte RE, Powell T, Freeman C. Radiological and pathological response following pre-operative radiotherapy for soft-tissue sarcoma. *Radiother Oncol.* 2010;97(3):404-407. doi:10.1016/j.radonc.2010.10.007
4. Le Grange F, Cassoni AM, Seddon BM. Tumour volume changes following pre-operative radiotherapy in borderline resectable limb and trunk soft tissue sarcoma. *Eur J Surg Oncol.* 2014;40(4):394-401. doi:10.1016/j.ejso.2014.01.011
5. Dangoor A, Seddon B, Gerrand C, Grimer R, Whelan J, Judson I. UK guidelines for the management of soft tissue sarcomas. *Clin Sarcoma Res.* 2016;6(1):20. doi:10.1186/s13569-016-0060-4
6. DAVIS A, OSULLIVAN B, TURCOTTE R, et al. Late radiation morbidity following randomization to preoperative versus postoperative radiotherapy in extremity soft tissue sarcoma. *Radiother Oncol.* 2005;75(1):48-53. doi:10.1016/j.radonc.2004.12.020

- 1 7. Eilber FC, Rosen G, Eckardt J, et al. Treatment-Induced Pathologic Necrosis: A
2 Predictor of Local Recurrence and Survival in Patients Receiving Neoadjuvant Therapy
3 for High-Grade Extremity Soft Tissue Sarcomas. Bönig H, ed. *J Clin Oncol*.
4 2001;19(13):3203-3209. doi:10.1200/JCO.2001.19.13.3203
5
6
7
8
- 9 8. Shah D, Borys D, Martinez SR, et al. Complete pathologic response to neoadjuvant
10 radiotherapy is predictive of oncological outcome in patients with soft tissue
11 sarcoma. *Anticancer Res*. 2012;32(9):3911-3915. doi:10.1038/mp.2011.182.doi
12
13
14
15
- 16 9. MacDermid DM, Miller LL, Peabody TD, et al. Primary Tumor Necrosis Predicts
17 Distant Control in Locally Advanced Soft-Tissue Sarcomas After Preoperative
18 Concurrent Chemoradiotherapy. Bönig H, ed. *Int J Radiat Oncol*. 2010;76(4):1147-
19 1153. doi:10.1016/j.ijrobp.2009.03.015
20
21
22
23
24
- 25 10. Miki Y, Ngan S, Clark JCMCM, Akiyama T, Choong PFMFM. The significance of size
26 change of soft tissue sarcoma during preoperative radiotherapy. *Eur J Surg Oncol*.
27 2010;36(7):678-683. doi:10.1016/j.ejso.2010.05.021
28
29
30
31
- 32 11. Pasquier D, Hadj Henni A, Escande A, et al. Diffusion weighted MRI as an early
33 predictor of tumor response to hypofractionated stereotactic boost for prostate
34 cancer. *Sci Rep*. 2018;8(1):10407. doi:10.1038/s41598-018-28817-9
35
36
37
38
- 39 12. Pham TT, Liney GP, Wong K, Barton MB. Functional MRI for quantitative treatment
40 response prediction in locally advanced rectal cancer. *Br J Radiol*.
41 2017;90(1072):20151078. doi:10.1259/bjr.20151078
42
43
44
45
- 46 13. Suh CH, Kim HS, Jung SC, Choi CG, Kim SJ. Multiparametric MRI as a potential
47 surrogate endpoint for decision-making in early treatment response following
48 concurrent chemoradiotherapy in patients with newly diagnosed glioblastoma: a
49 systematic review and meta-analysis. *Eur Radiol*. 2018;28(6):2628-2638.
50 doi:10.1007/s00330-017-5262-5
51
52
53
54
55
56
- 57 14. Chung PWM, Dehesi BM, Ferguson PC, et al. Radiosensitivity translates into
58 excellent local control in extremity myxoid liposarcoma: A comparison with other
59
60
61
62
63
64
65

- soft tissue sarcomas. *Cancer*. 2009;115(14):3254-3261. doi:10.1002/cncr.24375
15. Soldatos T, Ahlawat S, Montgomery E, Chalian M, Jacobs MA, Fayad LM. Multiparametric Assessment of Treatment Response in High-Grade Soft-Tissue Sarcomas with Anatomic and Functional MR Imaging Sequences. *Radiology*. 2016;278(3):831-840. doi:10.1148/radiol.2015142463
16. Meyer JM, Perlewitz KS, Hayden JB, et al. Phase I Trial of Preoperative Chemoradiation plus Sorafenib for High-Risk Extremity Soft Tissue Sarcomas with Dynamic Contrast-Enhanced MRI Correlates. *Clin Cancer Res*. 2013;19(24):6902-6911. doi:10.1158/1078-0432.CCR-13-1594
17. Xia W, Yan Z, Gao X. Volume fractions of DCE-MRI parameter as early predictor of histologic response in soft tissue sarcoma: A feasibility study. *Eur J Radiol*. 2017;95(July):228-235. doi:10.1016/j.ejrad.2017.08.021
18. Schnapauff D, Zeile M, Niederhagen M Ben, et al. Diffusion-weighted echo-planar magnetic resonance imaging for the assessment of tumor cellularity in patients with soft-tissue sarcomas. *J Magn Reson Imaging*. 2009;29(6):1355-1359. doi:10.1002/jmri.21755
19. O'Connor JPB, Aboagye EO, Adams JE, et al. Imaging biomarker roadmap for cancer studies. *Nat Rev Clin Oncol*. 2017;14(3):169-186. doi:10.1038/nrclinonc.2016.162
20. Maeda-Otsuka S, Kajihara I, Tasaki Y, et al. Hypoxia accelerates the progression of angiosarcoma through the regulation of angiosarcoma cells and tumor microenvironment. *J Dermatol Sci*. 2019. doi:10.1016/j.jdermsci.2019.01.005
21. Forker L, Gaunt P, Sioletic S, et al. The hypoxia marker CAIX is prognostic in the UK phase III Vortex-Biobank cohort: An important resource for translational research in soft tissue sarcoma. *Br J Cancer*. 2018. doi:10.1038/bjc.2017.430
22. Larsson HB, Rosenbaum S, Fritz-Hansen T. Quantification of the effect of water exchange in dynamic contrast MRI perfusion measurements in the brain and heart. *Magn Reson Med*. 2001;46(2):272-281.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

<http://www.ncbi.nlm.nih.gov/pubmed/11477630>.

23. Brookes JA, Redpath TW, Gilbert FJ, Murray AD, Staff RT. Accuracy of T1 measurement in dynamic contrast-enhanced breast MRI using two- and three-dimensional variable flip angle fast low-angle shot. *J Magn Reson Imaging*. 1999;9(2):163-171. <http://www.ncbi.nlm.nih.gov/pubmed/10077009>.
24. Yang L, Forker L, Irlam JJ, Pillay N, Choudhury A, West CML. Validation of a hypoxia related gene signature in multiple soft tissue sarcoma cohorts. *Oncotarget*. 2018;9(3):3946-3955. doi:10.18632/oncotarget.23280
25. Betts GNJ, Eustace A, Patiar S, et al. Prospective technical validation and assessment of intra-tumour heterogeneity of a low density array hypoxia gene profile in head and neck squamous cell carcinoma. *Eur J Cancer*. 2013;49(1):156-165. doi:10.1016/j.ejca.2012.07.028
26. McRobbie DW, Moore EA, Graves MJ, Prince MR. *MRI from Picture to Proton*. Cambridge: CUP; 2003.
27. Einarsdóttir H, Karlsson M, Wejde J, Bauer HCF. Diffusion-weighted MRI of soft tissue tumours. *Eur Radiol*. 2004;14(6):959-963. doi:10.1007/s00330-004-2237-0
28. Dudeck O, Zeile M, Pink D, et al. Diffusion-weighted magnetic resonance imaging allows monitoring of anticancer treatment effects in patients with soft-tissue sarcomas. *J Magn Reson Imaging*. 2008;27(5):1109-1113. doi:10.1002/jmri.21358
29. Jardim-Perassi B V., Huang S, Dominguez-Viqueira W, et al. Multiparametric MRI and coregistered histology identify tumor habitats in breast cancer mouse models. *Cancer Res*. 2019;79(15):3952-3964. doi:10.1158/0008-5472.CAN-19-0213
30. Surov A, Meyer HJ, Wienke A. Associations between apparent diffusion coefficient (ADC) and KI 67 in different tumors: A meta-analysis. part 2: ADCmin. *Oncotarget*. 2018;9(9):8675-8680. doi:10.18632/oncotarget.24006
31. Hompland T, Ellingsen C, Galappathi K, Rofstad EK. DW-MRI in assessment of the hypoxic fraction, interstitial fluid pressure, and metastatic propensity of melanoma

1 xenografts. *BMC Cancer*. 2014;14(1):1-9. doi:10.1186/1471-2407-14-92

- 2
3 32. Rasmussen JH, Olin A, Lelkaitis G, et al. Does multiparametric imaging with 18F-FDG-
4 PET/MRI capture spatial variation in immunohistochemical cancer biomarkers in head
5 and neck squamous cell carcinoma? *Br J Cancer*. 2020;123(1):46-53.
6 doi:10.1038/s41416-020-0876-9
7
8
9
10
11 33. Kerkmeijer LGW, Fuller CD, Verkooijen HM, et al. The MRI-Linear Accelerator
12 Consortium: Evidence-Based Clinical Introduction of an Innovation in Radiation
13 Oncology Connecting Researchers, Methodology, Data Collection, Quality Assurance,
14 and Technical Development. *Front Oncol*. 2016;6(October):1-6.
15 doi:10.3389/fonc.2016.00215
16
17
18
19
20
21
22
23 34. Nordsmark M, Alsner J, Keller J, et al. Hypoxia in human soft tissue sarcomas: Adverse
24 impact on survival and no association with p53 mutations. *Br J Cancer*.
25 2001;84(8):1070-1075. doi:10.1054/bjoc.2001.1728
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Tables

Table 1

Patient characteristics

Patient	Age at first scan	M/F	Tumour location	Tumour type, grade, stage	Status as of Jan 2019
1	73	M	Upper arm	Undifferentiated spindle cell G2, T2bN0M0	No disease
2	56	M	Trunk	Myxoid liposarcoma G3, T2bN0M0	No disease
3	79	M	Upper arm	Myxofibrosarcoma G2, T1bN0M0	No disease
4	27	M	Knee	Myxoid liposarcoma G3, T2bN0M0	No disease
5	29	M	Lower leg	Myxoid chondrosarcoma G3, T2bN0M0	No disease
6	69	F	Trunk	Undifferentiated pleomorphic sarcoma G3, T1aN0M0	No disease
7	41	M	Lower leg	Myxofibrosarcoma G3, T2bN0M0	No disease but chronic inflammation post-surgery
8	62	M	Forearm	Myxofibrosarcoma G3, T1bN0M0	No disease
9	24	M	Knee	Synovial sarcoma G3, T2bN0M0	Lung metastasis resected June 2018, now no disease
10	67	M	Knee	Undifferentiated pleomorphic sarcoma G3, T2bN0M0	Died (Acute Myeloid Leukaemia) Feb 2018
11	33	M	Thigh	Myxoid liposarcoma G3, T2bN0M0	Single metastasis in spine August 2018
12	74	M	Trunk	Undifferentiated spindle cell sarcoma G3, T2bN0M0	No disease

Table 2

Trial Imaging protocol

Sequence	Purpose	Flip angle, TE, TR / ms	Parallel imaging factor	Other	Matrix ^a	FOV / cm
TSE 2D Turbo spin- echo	High resolution T2w for tumour outlining	150° 96, 3890	None	ETL 13	256 x 256 x 20	Limb: 25 x 25 x 10 Trunk: 38.6 x 38.6 x 10
EPI 2D Echoplanar imaging	Diffusion- weighted imaging	- 103, 12100	2 AP	EPI factor 128, B = 0, 50, 100, 150, 200, 500, 1000 s/mm ²	128 x 128 x 20	
SRTFE 3D Saturation- recovery turbo field echo	T1 measurement	12° 1.52 64, 142, 292, 1050, 2550, 3950	2 AP	TI = 37, 100, 250, 1000, 2500, 3900 ms		
VIBE 3D Volume- interpolated breath hold imaging	Dynamic contrast- enhanced imaging	16° 0.81, 2.63	2 AP	Temporal resolution 1.75 s, 150 dynamics		

^a In one case, 26 slices were needed to cover the tumour, leading to a dynamic temporal resolution of 3.2 s, and TR values for the SRTFE of 73, 145, 306, 1060, 2560, and 3960 ms.

TE – echo time, TR – repetition time, TI – inversion time, FOV – field of view

Table 3

Pearson correlation coefficients for correlations between imaging and histological parameters., with p-values calculated from a t-distribution shown in brackets.

	T1	iAUC	ADC	Volume
	Pre-treatment			
CD31	-0.07 (0.83)	-0.28 (0.37)	-0.16 (0.63)	-0.37 (0.23)
Ki67	-0.50 (0.10)	0.11 (0.74)	-0.55 (0.06)	-0.29 (0.36)
HIF	0.08 (0.80)	-0.40 (0.19)	0.04 (0.91)	-0.10 (0.76)
Hypoxia score	0.47 (0.12)	-0.64 (0.03)*	0.63 (0.03)*	-0.16 (0.63)
	Mid-treatment			
CD31	-0.09 (0.78)	-0.43 (0.17)	-0.31 (0.32)	-0.32 (0.31)
Ki67	-0.43 (0.16)	-0.05 (0.88)	-0.66 (0.02)*	-0.31 (0.33)
HIF	0.03 (0.94)	-0.44 (0.16)	-0.10 (0.75)	-0.13 (0.68)
Hypoxia score	0.24 (0.45)	-0.63 (0.03)*	0.58 (0.05)	-0.14 (0.66)
	Post-treatment			
CD31	-0.15 (0.63)	-0.26 (0.42)	-0.29 (0.36)	-0.13 (0.69)
Ki67	-0.57 (0.05)	-0.09 (0.78)	-0.44 (0.16)	-0.04 (0.66)
HIF	-0.27 (0.40)	-0.20 (0.53)	-0.36 (0.25)	-0.29 (0.37)
Hypoxia score	0.23 (0.43)	-0.27 (0.40)	0.34 (0.27)	-0.18 (0.57)

Figure 1

Figure 1 - Modified CONSORT diagram




Figure 2 - Box plot showing median T1 (box middle line), lower and upper quartiles (box edges) and data range (whiskers) over all patients for three visits. Outliers are shown as diamonds. There was a significant difference between median T1 at visit 2 compared with visit 1 (Wilcoxon signed ranks test, $p=0.008$). No other significant differences were detected.




Figure 3 (colour) – Example ADC (apparent diffusion coefficient, $\times 10^{-6}$ mm²/s) maps superimposed on anatomical T2w images over three visits (left – before radiotherapy, centre – during radiotherapy, right – after radiotherapy) for two patients (Upper panel, lower panel).

